

JOURNAL OF ANDROLOGY

Volume 14

JANUARY/FEBRUARY 1993

Number 1

CONTENTS

Editorial	1
Androgen UDP-Glucuronyl Transferase Activity Is Found Primarily in Liver in the Rat / E. C. Pirog, R. V. Clark, D. C. Collins	2
Hormonal Modulation of the Interactions of Cultured Rat Testicular Sertoli and Peritubular Myoid Cells: Effects on Glycosaminoglycan Synthesis / S. S. Raychoudhury, A. W. Blackshaw, M. G. Irving	9
Decreased Responsiveness to Progesterone of Spermatozoa in Oligozoospermic Patients / C. Falsetti, E. Baldi, C. Krausz, R. Casano, P. Failli, G. Forti	17
Immunocytochemical Localization of the Y_r Subunit of Glutathione S-Transferase P Shows Regional Variation in the Staining of Epithelial Cells of the Testis, Efferent Ducts, and Epididymis of the Male Rat / J.-P. Veri, L. Hermo, B. Robaire	23
Pharmacokinetics and Pharmacodynamics of a Parenteral Testosterone Microsphere Formulation in the Male Rat: Demonstration of Dose Dependence and Controlled Release / N. R. Peacock, R. S. Swerdloff, N. Berman, R. M. Gilley, T. R. Tice, S. Bhasin	45
Importance of Total Motile Oval Count in Interpreting the Hamster Ovum Sperm Penetration Assay / V. T. Brandeis	53
Computer-Assisted Semen Analysis (CASA) of Epididymal Sperm from the Domestic Cat / J. J. Stachecki, K. A. Ginsburg, R. E. Leach, D. R. Armant	60
Effect of Catheter Composition on Sperm Quality / N. S. Davis, S. A. Rothmann, M. Tan, A. J. Thomas, Jr.	66
Announcements	American Society of Andrology
Notice to Copiers vi	Placement Service 8
North American Testis Workshop . . 16	Outstanding Original Research Award 22
	Annual Meetings 52

Cover: *Color confocal micrographs of a reconstructed area of the corpus epididymal epithelium immunostained for Y_r glutathione S-transferase P protein. This reconstruction was made up of 56 slices, each 0.3 μm thick. The reconstructed section is viewed at two different angles. Immunostaining intensity increases by approximately a factor of 10 between each color starting from blue, then green, red, and gray. In a cross-sectional view at 0° rotation (top), strong immunoreactivity (red, gray) is localized to the basal edge of the epithelium where basal cells reside, whereas very weak immunostaining (green, blue) is seen over principal cells or in the lumen. The extensive coverage of the tubule by the basal cells becomes more apparent as the tissue section is rotated through a 60° angle (bottom). × 1646. (See article by Veri et al)*