

**Temporal Decreases in Sperm Motility: Which Patients Should Have Motility Checked at  
Both One and Two Hours After Collection?**

Ryan T. Schulte<sup>1</sup>, Laura M. Keller<sup>2</sup>, Melissa R. Hiner<sup>2</sup>, Dana A. Ohl<sup>1</sup>, and  
Gary D. Smith<sup>1,2,3,4,5</sup>

Departments of Urology<sup>1</sup>, Obstetrics and Gynecology<sup>2</sup>, Integrated and Molecular Physiology<sup>3</sup>,  
and Reproductive Sciences Program<sup>4</sup>, University of Michigan, Ann Arbor, MI 48109-0617

<sup>5</sup>Correspondence: Gary D. Smith, PhD  
6428 Medical Sciences I  
1301 E. Catherine St  
Ann Arbor, MI 48109-0617  
Tel: (734) 764-4134  
Fax (734) 936-8617  
Email: smithgd@umich.edu

Running Title: Temporal Decreases in Sperm Motility

**Abstract:**

A decrease in sperm motility and thus total motile sperm count (TMSC) over a period of hours may have clinical implications in counseling couples considering intrauterine insemination (IUI), *in vitro* fertilization (IVF), and intracytoplasmic sperm injection (ICSI). The objective of this study was to identify patients with decreases in sperm motility from 1 hour to 2 hours after collection and examine predictive relationships with semen analysis parameters. Between 2001 and 2005, 2313 semen samples were analyzed. Sperm motility was evaluated at both 1 and 2 hours after time of collection. Relevant seminal parameters were compared between patients with a decrease from the 1-hour to 2-hour motility (n=384) vs. those that showed no change (n=1929). The same analysis was performed in a subset of patients with a TMSC between 10-40 million. In the total patient population, only 16% (384/2313) demonstrated a decrease in the 1-hour to 2-hour motility. In patients displaying a decrease in the 1-2 hour motility, sperm concentration (33.5 vs. 79 mil/ml,  $p < 0.0001$ ) and % normal morphology (7% vs. 8%,  $p < 0.0001$ ) were significantly lower. Additionally, a significantly higher incidence of 1-2 hour motility decrease was seen in patients with midpiece anomalies (33.3% vs. 15.9%,  $p = 0.01$ ). Within the subset population of TMSC 10-40 million, the only statistically significant difference was in patients with midpiece anomalies (80.0% vs. 28.2%,  $p = 0.02$ ) who demonstrated a higher incidence of 1-2 hour motility decrease. Overall, patients with a TMSC between 10-40 million showed a significantly higher incidence of 1-hour to 2-hour motility decrease compared to the rest of the patient population (29.0% vs. 14.6%,  $p < 0.0001$ ). Because decreases in the 1 to 2-hour sperm motility affect only a small portion of patients, it is not necessary to check 2-hour motility on all patients. However, because patients with a TMSC between 10-40 million were significantly more likely to show a decrease in sperm motility, a decrease which could have possible clinical implications in couples deciding between IUI, IVF or ICSI, checking 2-hour sperm motility should be considered in this population.

**Keywords:** Assisted reproduction; Infertility; Semen; Semen Analysis

## Introduction

Male factor infertility is the sole cause of infertility in approximately 20% of infertile couples, and is a contributing factor in another 30% to 40% (Mosher and Pratt 1991; Thonneau et al, 1991). Thus, semen analyses remain an essential component of infertility evaluation. Normal semen quality is usually based on World Health Organization (WHO) criteria for multiple seminal parameters including seminal volume, pH, sperm concentration, motility, and morphology (World Health Organization 1999). The WHO criteria for normality of sperm motility is 50% or more with forward progression within 60 minutes of ejaculation (World Health Organization 1999). Given these criteria, many laboratories may not check sperm motility after more than 1 hour.

Sperm motility is an important factor to consider in couples pursuing assisted reproductive technologies (ART). Several studies have demonstrated the influence of sperm motility on the outcomes of various ARTs, including intrauterine insemination (IUI) (Lee et al, 2002; Yalti et al, 2004; Zhao et al, 2004), conventional *in vitro* fertilization (IVF) (Donnelly et al, 1998; Repping et al, 2002), and IVF with intracytoplasmic sperm injection (ICSI) (Stalf et al, 2005). Although the exact TMSC cutoff for defining severe male factor infertility varies somewhat among studies, a TMSC of 10 million in the ejaculate has been reported to be a useful threshold for decisions about treating a couple with IUI or IVF (Van Voorhis et al, 2001). This is also many times a practical threshold for conventional insemination or ICSI. Thus, a decrease in sperm motility and thus TMSC over a period of a few hours could have relevant clinical implications in counseling these couples whether to pursue IUI, IVF or ICSI. This decrease could have an even greater effect in oligospermic patients with already borderline low TMSC around this threshold value of 10 million.

The objective of this study was to determine the incidence of decreases in sperm motility seen from 1 hour to 2 hours after collection in a large population of patients presenting to an infertility clinic and to identify related seminal parameters which help predict these decreases. In this retrospective study over a four year period, all patients who demonstrated a decrease in sperm motility from 1 to 2 hours after collection were identified and multiple relevant seminal parameters were analyzed to identify any predictive parameters. A similar subset analysis was also performed on patients with a baseline TMSC between 10-40 million, as decreases in this population are more likely to cause the TMSC to drop below the aforementioned threshold value

of 10 million, and thus, be useful for decision making in couples contemplating IUI, IVF, or ICSI.

## **Methods**

### **Semen Analysis**

Use of semen analysis outcome data for investigative purposes was approved following Investigational Review Board (IRB) application and review. Semen samples were collected from men presenting to the University of Michigan Center for Reproductive Medicine. All men were instructed to refrain from any sexual activity including masturbation for approximately 3-5 days prior to sample collection. Semen was collected by masturbation in a lab-approved sterile plastic container. Semen specimens were allowed to liquefy at 37° C prior to analysis approximately one hour after ejaculation. Semen volume was measured with a graduated pipette to the nearest 0.1 ml. A drop of semen was placed on pHDrion paper (Micro Essential Laboratory, Inc., Brooklyn, NY) and pH was measured and recorded. Sperm concentration was determined for duplicate loadings on a Neubauer hemocytometer. Sperm motility was assessed at both 1 and 2 hours after collection. For assessment of motility, a minimum of 200 sperm from multiple fields were counted using the 37° C warming stage and 20x magnification. Percentage of motile sperm with forward progression and speed of progression were also analyzed at this time. For assessment of morphology, 10 µl of semen was smeared and air-dried, then stained with Papanicolau stain. Morphologic assessment was then carried out under the 100X objective with immersion oil until 200 sperm were counted. Sperm morphology was determined using the strict criteria described by Kruger et al (1986) at 100X magnification. These gross head defects, mid-piece defects, and tail defects also were scored. Presence of anti-sperm antibodies was evaluated using Sperm-Mar (MAR; Fertipro NV, Belgium).

### **Statistical Analysis**

Information was collected retrospectively on 2313 semen analyses collected from 2003 to 2006. Patients were divided into two groups: 1) those that showed a decrease in motility from 1 hour to 2 hours after collection (n=384), and 2) those that showed no change in motility from 1 hour to 2 hours after collection (n=1929). Decrease in motility was defined as those samples that had a  $\geq 10\%$  decline in the 2 hour motility when compared to the 1 hour motility. Statistical

analysis using Fisher's exact test was performed to evaluate the bivariate relationship between a decrease in motility from 1 hour to 2 hours after collection and other categorical semen analysis variables. The Wilcoxon two-sample test was used to compare the bivariate relationship between a decrease in motility from 1 hour to 2 hours after collection and continuous semen analysis variables. A P-value less than 0.05 was considered statistically significant.

Additionally, a similar subset analysis was performed on patients with a baseline TMSC between 10-40 million (n=321). Among this subset, patients were again divided into two groups: 1) those that showed a decrease in motility from 1 hour to 2 hours after collection (n=93), and 2) those that showed no change in motility from 1 hour to 2 hours after collection (n=228). Fisher's exact test and Wilcoxon two-sample test were again used to evaluate the bivariate relationship between a decrease in motility from 1 hour to 2 hours after collection with categorical and continuous semen analysis variables, respectively.

## Results

In the total patient population, only 16% of samples (384/2313) demonstrated a decrease in the 1-hour to 2-hour motility. Table 1 shows the comparison of relevant seminal parameters between the group that demonstrated a decrease in the 1-hour to 2-hour motility versus those with no change. The range of sperm concentrations in the entire data set was 0 to  $2,432 \times 10^6$ . Sperm concentration ( $p < 0.0001$ ), percent normal morphology ( $p < 0.0001$ ), and percent of motile sperm with forward progression ( $p = 0.0001$ ) were all lower in patients that had a decrease in the 1-hour to 2-hour motility when compared to patients with no change in motility. Additionally, the percentage of abnormal sperm forms with head defects ( $p = 0.001$ ) and midpiece defects as well as the percentage of sperm bound with IgG anti-sperm antibodies ( $p = 0.01$ ) were higher in patients that had a decrease in the 1-hour to 2-hour motility. Table 2 displays several other categorical seminal parameters that were evaluated between the two groups. A greater incidence of 1-hour to 2-hour motility decrease was seen in patients whose semen analysis had morphologic midpiece defects ( $p = 0.01$ ), greater than  $5 \times 10^6$  round cells ( $p = 0.01$ ), incomplete liquefaction ( $p = 0.04$ ), and a sluggish speed of progression ( $p < 0.0001$ ).

The TMSC range for the total population was 0 to 967 million. However, in our subset analysis, patients with a TMSC between 10-40 million were almost twice as likely to have a drop in motility from 1-hour to 2-hours after collection compared to the rest of the population (29.0%

vs. 14.6%,  $p < 0.0001$ ; Figure 1). The results of the subset analysis of patients with a TMSC between 10-40 million are shown in Tables 1 and 2. The only statistically significant difference among the multiple seminal parameters examined was seen in patients with gross morphologic midpiece defects, who subsequently had a higher incidence of 1-hour to 2-hour motility decrease compared to those without midpiece defects ( $p = 0.03$ ). Among both the total patient population and the subpopulation of TMSC between 10-40 million, the presence of midpiece defects was the only seminal parameter that showed a statistically significant relationship to the incidence of 1-hour to 2-hour motility decrease in both populations (Figure 2).

## Discussion

Semen analysis is an integral component of evaluation of infertility. Sperm motility is an important factor in the fertilizing potential of the sperm, and motility has been shown to correlate closely with fertilization rates of human oocytes *in vitro* (Bongso et al, 1989). Subsequently, sperm motility is also an important factor in the outcomes of various ARTs. Multiple studies have shown that IUI outcomes are affected by the percentage of motile sperm and total motile sperm count (Shulman et al, 1998; Pasqualotto et al, 1999; Lee et al, 2002; Miller et al, 2002; Yalti et al, 2004; Zhao et al, 2004). This effect on IUI outcomes has been demonstrated with both the percentage of motile sperm in the unprocessed ejaculate (Lee et al, 2002; Yalti et al, 2004; Zhao et al, 2004) as well as the post-processing sperm motility (Shulman et al, 1998; Pasqualotto et al, 1999). Both percentage of motile sperm and TMSC in unprocessed semen have also been shown to influence IVF outcomes (Donnelly et al, 1998; Van Voorhis et al, 2001; Repping et al, 2002). Additionally, the fertilization potential of injected sperm during ICSI was found to also be associated with motility in both ejaculated and testicular sperm (Stalf et al, 2005). Given this impact of sperm motility on ART outcomes, a decline in motility over a period of a few hours could have relevant clinical implications in couples pursuing ARTs.

Many laboratories do not routinely check repeated measures of sperm motility. Sperm motility can be time dependent. Under normal laboratory conditions, sperm motility will gradually decline, although significant signs of change are not usually apparent until three hours have elapsed (Glover et al, 1990). A decrease in motility of up to 20% in 3-4 hours is not unusual, but if the fall-off is greater or quicker, infection or some other abnormality should be suspected (Glover et al, 1990). Previous studies have looked at sperm motility longevity. Denil

and coworkers (1992) reported that after 24 hours, samples from men with male factor infertility lost 48% of initial motility while normal specimens lost only 34%. Makler and colleagues (1979) found three typical curves of motility change during the first 4 hours: increase in motility, moderate decline in motility, and rapid loss of motility. It was in this population of patients with rapid loss of motility over an hour that we sought to identify the incidence and evaluate their relationship with other relevant seminal parameters. In our retrospective study of 2313 semen analysis over a four year period, we found that decreases in sperm motility from 1-hour to 2-hours after collection affected only a relatively small proportion of patients, with an incidence of 16%. Singer and coworkers (1980) reported that percentile decreases in sperm motility after 5 hours were found to be higher in oligospermic specimens than in those with higher sperm densities. In the present study, patients with decreases in motility from 1-hour to 2-hours after collection demonstrated lower mean sperm concentrations when compared to patients that showed no change in sperm motility. Although this difference was statistically significant, the mean sperm concentration in both groups was still high enough to be considered normospermic and thus would likely be of little clinical significance in such a way that would help guide which patients should have motility checked at both 1-hour and 2-hours post-collection. Percent normal morphology and percent of motile sperm with forward progression were also lower in patients with decreases in 1-hour to 2-hour sperm motility. Similarly, although a significant observation from a statistical perspective, the actual differences are too small to affect clinical decision making in determining which patients should have sperm motility checked at both 1-hour and 2-hours.

Because these decreases in sperm motility over a period of a couple hours affect only a small portion of patients, we propose that it is unnecessary to check both 1-hour and 2-hour motility on all patients. Even in some patients who display a decrease in motility, ART clinical decision making may be unaffected. For example, in a couple contemplating ART, if the man already has a high TMSC over 100 million, even after a 50% decrease in sperm motility, he would still have a relatively high TMSC, and IUI would still be the likely be the recommended first line ART for the couple. On the other end of the spectrum, in a man with severely low TMSC of less than 1 million, a 50% decrease in sperm motility would also likely not affect the ART decision making, as ICSI would likely be recommended either way. However, the patients that may be most affected by a decrease in sperm motility would be those with a TMSC in

between these two extremes. The exact TMSC threshold for defining severe male factor infertility has varied somewhat among studies, but threshold values of between 5 and 10 million in the ejaculate have been reported (Cohlen et al, 1998; Dickey et al, 1999; Van Voorhis et al, 2001). Van Voorhis and colleagues (2001) reported that the average TMSC in the ejaculate was a useful independent predictor of clinical pregnancy after IUI, with a threshold value of 10 million. Additionally, when the TMSC in the ejaculate was under 10 million, IVF with ICSI was more cost-effective than IUI. A decrease in sperm motility and TMSC over a period of an hour would have the most relevant clinical implications in patients with a moderately low TMSC around this 10 million threshold and the subsequent decision whether to pursue IUI, IVF with conventional insemination or ICSI. Thus, we performed a subset analysis in this population, identifying patients with a TMSC between 10-40 million. Patients in this subpopulation were twice as likely to show a decrease in the 1-hour to 2-hour motility when compared to the rest of the population (29.0% vs. 14.6%,  $p < 0.0001$ ). Given the significantly increased incidence of motility decrease in patients with a TMSC between 10-40 million and its subsequent potential clinical implications for couples deciding between various ART interventions, checking both 1-hour and 2-hour sperm motility should be considered in this population.

Interestingly, the only common seminal parameter that showed a statistically significant relationship to the incidence of 1-hour to 2-hour motility decrease in both the total patient population and the subpopulation of TMSC between 10-40 million was presence of gross morphologic midpiece defects. The sperm midpiece is densely packed with mitochondria which provide energy in the form of ATP to the flagellum to propel the cell. Multiple studies have examined the relationship between the sperm mid-piece and mitochondria with sperm motility. Motility depends largely on the mitochondrial volume within the sperm midpiece (Ruiz-Pesini et al, 1998) and morphologic defects of the midpiece can cause non-progressive movement or immotility (Piasecka and Kawiak 2003). Mundy and colleagues (1995) showed that sperm mid-piece length was significantly shorter in asthenozoospermic subjects compared with controls. Additionally, Nakada and coworkers (2006) used a transmittochondrial mouse model to show that accumulation of mutant mitochondrial DNA induces asthenozoospermia with the majority of sperm showing midpiece abnormalities. In the present study, patients with morphologic defects of sperm midpiece on semen analysis were more than twice as likely to show a decrease in the 1-hour to 2-hour sperm motility when compared with those with no midpiece defects. This

relationship was seen consistently across our total patient population as well as in our subpopulation analysis. The current study is subject to the limitations of light microscopy to evaluate morphologic sperm defects. Further studies with electron microscopy would be necessary to fully evaluate to the true morphologic structural abnormalities contributing to these temporal decreases in motility. However, given our current findings in combination with the multiple previous studies, checking both 1-hour and 2-hour sperm motility should also be considered in patients with morphological midpiece defects on semen analysis.

So far as is known, this is the first study to determine the incidence and related seminal parameters of rapid declines in sperm motility longevity over a 1 hour period in a such a large sample of subjects from an infertility clinic population. In conclusion, the incidence of rapid declines in sperm motility from 1 hour to 2 hours after collection is relatively low, affecting only 16% of the population. Thus, it is likely unnecessary to make 2-hour sperm motility a routine part of semen analysis for all patients. However, patients with a TMSC between 10-40 million, as well as patients with morphologic midpiece defects of the sperm, had a significantly higher incidence of rapid declines in sperm motility over the 1 hour period. Because of the possible ART clinical implications of rapid declines in sperm motility in these patients, checking both 1-hour and 2-hour sperm motility should be considered in these populations.

### **Acknowledgments**

The authors would like to thank Dr. Carrie Cosola-Smith for critical review of this manuscript and editorial comments. The authors would also like to thank Rodney L. Dunn, MS for his assistance with statistical analysis.

### **References**

Bongso TA, Ng SC, Mok H, Lim MN, Teo HL, Wong PC, Ratnam SS. Effect of sperm motility on human in vitro fertilization. *Arch Androl.* 1989;22:185-190.

Cohlen BJ, te Velde ER, van Kooij RJ, Looman CW, Habbema JD. Controlled ovarian hyperstimulation and intrauterine insemination for treating male subfertility: a controlled study. *Hum Reprod.* 1998;13:1553-1558.

Denil J, Ohl DA, Hurd WW, Menge AC, Hiner MR. Motility longevity of sperm samples processed for intrauterine insemination. *Fertil Steril.* 1992;58:436-438.

- Dickey RP, Pyrzak R, Lu PY, Taylor SN, Rye PH. Comparison of the sperm quality necessary for successful intrauterine insemination with World Health Organization threshold values for normal sperm. *Fertil Steril*. 1999;71:684-689.
- Donnelly ET, Lewis SE, McNally JA, Thompson W. In vitro fertilization and pregnancy rates: the influence of sperm motility and morphology on IVF outcome. *Fertil Steril*. 1998;70:305-314.
- Glover T, Barrat C, Tyler J, Hennessey J. *Human male fertility and semen analysis*. London: Academic Press Limited; 1990.
- Kruger TF, Menkveld R, Stander FS, Lombard CJ, Van der Merwe JP, van Zyl JA, Smith K. Sperm morphologic features as a prognostic factor in in vitro fertilization. *Fertil Steril*. 1986;46:1118-1123.
- Lee VM, Wong JS, Loh SK, Leong NK. Sperm motility in the semen analysis affects the outcome of superovulation intrauterine insemination in the treatment of infertile Asian couples with male factor infertility. *Bjog*. 2002;109:115-120.
- Makler A, Zaidise I, Paldi E, Brandes JM. Factors affecting sperm motility. I. In vitro change in motility with time after ejaculation. *Fertil Steril*. 1979;31:147-154.
- Miller DC, Hollenbeck BK, Smith GD, Randolph JF, Christman GM, Smith YR, Lebovic DI, Ohl DA. Processed total motile sperm count correlates with pregnancy outcome after intrauterine insemination. *Urology*. 2002;60:497-501.
- Mosher WD, Pratt WF. Fecundity and infertility in the United States: incidence and trends. *Fertil Steril*. 1991;56:192-193.
- Mundy AJ, Ryder TA, Edmonds DK. Asthenozoospermia and the human sperm mid-piece. *Hum Reprod*. 1995;10:116-119.
- Nakada K, Sato A, Yoshida K, Morita T, Tanaka H, Inoue S, Yonekawa H, Hayashi J. Mitochondria-related male infertility. *Proc Natl Acad Sci U S A*. 2006;103:15148-15153.
- Pasqualotto EB, Daitch JA, Hendin BN, Falcone T, Thomas AJ, Jr., Nelson DR, Agarwal A. Relationship of total motile sperm count and percentage motile sperm to successful pregnancy rates following intrauterine insemination. *J Assist Reprod Genet*. 1999;16:476-482.
- Piasecka M, Kawiak J. Sperm mitochondria of patients with normal sperm motility and with asthenozoospermia: morphological and functional study. *Folia Histochem Cytobiol*. 2003;41:125-139.
- Repping S, van Weert JM, Mol BW, de Vries JW, van der Veen F. Use of the total motile sperm count to predict total fertilization failure in in vitro fertilization. *Fertil Steril*. 2002;78:22-28.

Ruiz-Pesini E, Diez C, Lapena AC, Perez-Martos A, Montoya J, Alvarez E, Arenas J, Lopez-Perez MJ. Correlation of sperm motility with mitochondrial enzymatic activities. *Clin Chem*. 1998;44:1616-1620.

Shulman A, Hauser R, Lipitz S, Frenkel Y, Dor J, Bider D, Mashiach S, Yogev L, Yavetz H. Sperm motility is a major determinant of pregnancy outcome following intrauterine insemination. *J Assist Reprod Genet*. 1998;15:381-385.

Singer R, Sagiv M, Landau B, Allalouf D, Segenreich E, Servadio C. Motility and vitality of human spermatozoa at various time intervals after ejaculation. *Experientia*. 1980;36:578-579.

Stalf T, Mehnert C, Hajimohammad A, Manolopoulos K, Shen Y, Schuppe HC, Diemer T, Schill WB, Weidner W, Tinneberg HR. Influence of motility and vitality in intracytoplasmic sperm injection with ejaculated and testicular sperm. *Andrologia*. 2005;37:125-130.

Thonneau P, Marchand S, Tallec A, Ferial ML, Ducot B, Lansac J, Lopes P, Tabaste JM, Spira A. Incidence and main causes of infertility in a resident population (1,850,000) of three French regions (1988-1989). *Hum Reprod*. 1991;6:811-816.

Van Voorhis BJ, Barnett M, Sparks AE, Syrop CH, Rosenthal G, Dawson J. Effect of the total motile sperm count on the efficacy and cost-effectiveness of intrauterine insemination and in vitro fertilization. *Fertil Steril*. 2001;75:661-668.

World Health Organization. *WHO Laboratory Manual for the Examination of Human Semen and Sperm-Cervical Mucus Interaction*. 4th edition. Cambridge, United Kingdom: Cambridge University Press; 1999.

Yalti S, Gurbuz B, Sezer H, Celik S. Effects of semen characteristics on IUI combined with mild ovarian stimulation. *Arch Androl*. 2004;50:239-246.

Zhao Y, Vlahos N, Wyncott D, Petrella C, Garcia J, Zacur H, Wallach EE. Impact of semen characteristics on the success of intrauterine insemination. *J Assist Reprod Genet*. 2004;21:143-148.

**TABLE 1.** Seminal parameters in total patient population and subpopulation of total motile sperm count  $10-40 \times 10^6$ .

Semen Parameter	Total patient population			Subpopulation: TMSC <sup>‡</sup> 10-40 x10 <sup>6</sup>		
	1-2h Motility Decrease	No Motility Change	P-Value	1-2h Motility Decrease	No Motility Change	P-Value
Days of abstinence	4.8 ± 5.4	4.7 ± 6.2	0.35	5.0 ± 9.1	5.4 ± 15.3	0.56
Volume (mL)	3.3 ± 1.9	3.3 ± 1.6	0.51	3.0 ± 1.9	2.6 ± 1.6	0.10
pH	7.53 ± 0.4	7.55 ± 0.2	0.82	7.49 ± 0.6	7.57 ± 0.2	0.21
Concentration (x10 <sup>6</sup> /mL)	64.4 ± 109.1	102.5 ± 109.9	<0.0001*	38.8 ± 51.2	35.3 ± 37.1	0.77
Forward progression (%) <sup>#</sup>	69.6 ± 15.0	72.5 ± 13.2	0.0001*	67.7 ± 14.9	71.3 ± 6.3	0.44
Morphology (% normal forms)	7.4 ± 4.0	8.6 ± 4.8	<0.0001*	7.5 ± 3.4	7.1 ± 2.9	0.29
Head anomalies (% of abnormal forms)	90.6 ± 4.7	89.6 ± 5.7	0.001*	89.8 ± 4.1	90.2 ± 7.1	0.06
Midpiece defects (% of abnormal forms)	0.08 ± 0.5	0.02 ± 0.2	0.005*	0.05 ± 0.2	0.004 ± 0.06	0.01*
Tail anomalies (% of abnormal forms)	1.75 ± 3.1	1.87 ± 2.4	0.22	2.47 ± 2.6	2.63 ± 6.3	0.64
IgA anti-sperm Ab (% sperm bound)	5.8 ± 19.2	2.6 ± 9.2	0.52	8.0 ± 25.4	3.3 ± 11.1	0.25
IgG anti-sperm Ab (% sperm bound)	8.4 ± 25.6	3.9 ± 17.3	0.01*	12.0 ± 30.5	6.2 ± 22.4	0.24

<sup>‡</sup> Total motile sperm count

<sup>#</sup> % of motile sperm with forward progression

\* Statistically significant

**TABLE 2.** Categorical seminal parameters in total patient population and subpopulation of total motile sperm count  $10-40 \times 10^6$ .

<b>Semen Parameter</b>	<b>Total Patient Population</b>		<b>Subpopulation: TMS<sup>‡</sup> 10-40x10<sup>6</sup></b>	
	<b>Incidence of 1-2h Motility Decrease (%)</b>	<b>P-Value</b>	<b>Incidence of 1-2h Motility Decrease (%)</b>	<b>P-Value</b>
Place of Collection				
Home	16.6	0.90	30.1	0.88
Lab	16.8		28.6	
Liquefaction				
Complete	16.5	0.04*	28.8	0.29
Incomplete	38.5		100.0	
Agglutination				
None	22.5	0.13	31.1	0.14
Slight	12.4		25.8	
Moderate	16.7		100.0	
Heavy	40.0		100.0	
Speed of Progression				
None-sluggish	24.6	<0.0001*	33.3	0.24
Good-excellent	14.3		26.8	
Round Cells				
<5x10 <sup>6</sup> /mL	16.2	0.01*	28.5	0.29
≥5x10 <sup>6</sup> /mL	28.8		44.4	
Midpiece anomaly				
Absent	15.9	0.01*	28.2	0.01*
Present	33.3		80.0	

<sup>‡</sup> Total motile sperm count


\* Statistically significant

**Figure Legends**

**Figure 1.** Incidence of decreases in 1-hour to 2-hour sperm motility in patients with total motile sperm count (TMSC)  $10-40 \times 10^6$  compared to the remainder of the patient population.

**Figure 2.** Relationship of sperm morphologic midpiece defects to the incidence of decreases in 1-hour to 2-hour sperm motility

Figure 1.



**Figure 2.**